

ACKNOWLEDGMENTS

We thank the many persons who assisted with this study. Those who aided in organization and planning were: Herbert W. Graham, Robert L. Edwards, and K. O. Emery. Personnel from the Northeast Fisheries Center who assisted with the biological work were: Bruce R. Burns, Philip H. Chase, Jr., Christopher P. Ciano, Evan B. Haynes, Henry W. Jensen, Barbara Langevin, Lewis M. Lawday, Arthur S. Merrill, Harriett E. Murray, Clifford D. Newell, Carol Schwamb, Ruth Stoddard Byron, Rodman E. Taylor, Jr., and Ellen J. Winchell. James O. Farlow, Edward M. Handy, and John P. Laird provided assistance with computer programming. Frank A. Bailey and John R. Lamont assisted with drafting. Scientists from the U.S. Geological Survey and Woods Hole Oceanographic Institution marine geology group who provided sedimentological information or participated in shipboard work, of which biological sampling was an ancillary part, were: John C. Hathaway, Jobst Hülsemann, Frank Manheim, Robert H. Meade, Richard M. Pratt, David Ross, John S. Schlee, James V. A. Trumbull, and Elazar Uchupi.

Those who generously provided taxonomic assistance were: Edward L. Bousfield, Susan Brown-Leger, John C. McCain, Edward B. Cutler, Lion F. Gardner, Porter M. Kier, Peter Kinner, Louis S. Kornicker, John N. Kraeuter, Don Maurer, Arthur S. Merrill, Roy Oleröd, David L. Pawson, Frank Perron, Marian H. Pettibone, Thomas PheJan, Harold H. Plough, Johanna Reinhart, Howard L. Sanders, Thomas J. M. Schopf, Eve C. Southward, J. H. Stock, Lowell P. Thomas, Ruth D. Turner, Bernt Widersten, Austin B. Williams, Lev A. Zenkevitch, and Victor A. Zullo.

Captain Arthur D. Colburn, Jr. of R/V Asterias, Captain Walter E. Beatteay and crew of R/V Albatross IV, Captain John J. Walsh and crew of R/V Delaware I, and Captain Harry Seibert and crew of R/V Gosnold assisted with the collection of biological samples and physical-chemical data at sea.

Coordination with the Marine Ecosystems Analysis Program, New York Bight Project, was handled by Joel O'Connor and R. L. Swanson.

LITERATURE CITED

- Adams, C.B., 1839, Observations on some species of the marine shells of Massachusetts, with descriptions of five new species: Boston Jour. Natural History, v. 2, p. 262-288.
- Agassiz, E.C., and Agassiz, Alexander, 1865, Seaside studies in natural history. Marine animals of Massachusetts Bay. Radiates: Boston, Ticknor and Fields, 155 p.
- Aller, B.B., 1958, Publications of the United States Bureau of Fisheries 1871-1940: U.S. Dept. of the Interior, Fish and Wildlife Service, Spec. Sci. Rept. - Fisheries No. 284, 202 p.
- Anonymous, 1974, Living resource - related aspects of ocean dumping in the New York Bight: U.S. Dept. Commerce, NOAA, NMFS, Middle Atlantic Coastal Fisheries Center, Highlands, N.J., 07732, unpub. document, 15 p. and Appendix with 4 tables and 34 figs.
- Ayers, J.C., 1951, A preliminary report upon the state of our knowledge of the waters in and around New York Harbor, Appendix B, Bibliography, in Ayers, John C., et al., The Hydrography of the New York Area: Contract N6 DNR 264, task 15, Cornell University, Ithaca, New York.
- Barnes, R.D., 1974, Invertebrate zoology. Third edition: Philadelphia, London, Toronto, W.B. Saunders Company, 870 p.
- Bigelow, H.B., 1933, Studies of the waters on the continental shelf, Cape Cod to Chesapeake Bay; I. The cycle of temperature: Massachusetts Inst. Technology and Woods Hole Oceanog. Inst., Papers in Physics, Oceanography, and Meteorology, v. 2, no. 4, p. 1-135.

- Bigelow, H.B., and Sears, Mary, 1935, Studies of the waters of the continental shelf, Cape Cod to Chesapeake Bay: II, Salinity: Massachusetts Inst. Technology and Woods Hole Oceanog. Inst., Papers in Physical Oceanography and Meteorology, v. 4, no. 1, 94 p.
- Boesch, D.F., 1972, Species diversity of marine macrobenthos in the Virginia area: Ches. Sci., v. 13, p. 206-211.
- \_\_\_\_\_ 1973, Classification and community structure of macrobenthos in the Hampton Roads area, Virginia: Marine Biology, v. 21, p. 226-244.
- Bumpus, D.F., 1965, Residual drift along the bottom on the continental shelf in the Middle Atlantic Bight area: Limnology and Oceanography, v. 10, Supp., p. R50-R53.
- Bumpus, D.F., and Lauzier, L.M., 1965, Surface circulation on the continental shelf off eastern North America between Newfoundland and Florida: Am. Geog. Soc., Serial Atlas Marine Environment, Folio 7.
- Bumpus, D.F., Lynde, R.E., and Shaw, D.M., 1973, Physical oceanography, in Coastal and offshore environmental inventory, Cape Hatteras to Nantucket Shoals: Univ. Rhode Island, Marine Publication Series No. 2, Occasional Publ., no. 5, p. 1-1 to 1-72.
- Burbanck, W.D., Pierce, M.E., and Whiteley, G.C., Jr., 1956, A study of the bottom fauna of Rand's Harbor, Massachusetts: An application of the ecotone concept: Ecological Monographs, v. 26, no. 3, p. 213-243.
- Chase, Joseph, 1959, Wind-induced changes in the water column along the East Coast of the United States: Jour. Geophys. Research, v. 64, no. 8, p. 1013-1022.

- Churgin, James, and Halminski, S.J., 1974, Temperature, salinity, oxygen, and phosphate in waters off United States. Volume 1. Western North Atlantic: Key to Oceanographic Records Documentation No. 2, NODC, NOAA, 1-166 p.
- Colton, J.B., Jr., and Stoddard, R.R., 1972, Average monthly sea-water temperatures, Nova Scotia to Long Island, 1940-1959: Am. Geog. Soc., Serial Atlas Marine Environment, Folio 21.
- \_\_\_\_\_ 1973, Bottom-water temperatures on the continental shelf, Nova Scotia to New Jersey: NOAA Technical Rept., NMFS CIRC - 376, 55 p.
- D'Agostino, A., and Colgate, W., 1973, The Benthic organisms of the New York Bight, September and November 1971: a limited study of species and sediments: New York Ocean Science Lab. Tech. Rept. No. 0017, v. 2, p. 189-205.
- Dean, David, 1975, Raritan Bay macrobenthos survey, 1957-1960: National Oceanic and Atmospheric Administration, (NOAA), National Marine Fisheries Service (NMFS), NMFS Data Rept. 99, 51 p.
- Dean, David, and Haskin, H.H., 1964, Benthic repopulation of the Raritan River Estuary following pollution abatement: Limnology and Oceanography, v. 9, no. 4, p. 551-563.
- Desor, E., 1851, On echinoderms: Boston Soc. Nat. Hist. Proc., v. 3, (1848-1851), p. 65-68.
- Emery, K.O., 1966, The Atlantic Continental Shelf and Slope of the United States-geologic background: U.S. Geol. Survey Prof. Paper 529-A, 23 p.

- \_\_\_\_\_ 1968, Relict sediments on continental shelves of the world: Am. Assoc. Petroleum Geologists Bull., v. 52, no. 3, p. 445-464.
- Emery, K.O., and Merrill, A.S., 1964, Combination camera and bottom grab: Oceanus, v. 10, no. 4, p. 2-5.
- Emery, K.O., Merrill, A.S., and Trumbull, J.V.A., 1965, Geology and biology of the sea floor as deduced from simultaneous photographs and samples: Limnology and Oceanography, v. 10, no. 1, p. 1-21.
- Emery, K.O., and Schlee, J.S., 1963, The Atlantic continental shelf and slope, a program for study: U.S. Geol. Survey Circ. 481, 11 p.
- Emery, K.O., and Uchupi, Elazar, 1965, Structure of Georges Bank: Marine Geology, v. 3, p. 349-358.
- \_\_\_\_\_ 1972, Western North Atlantic Ocean: Topography, rocks, structure, water, life, and sediments: Am. Assoc. Petroleum Geologists Mem. 17, 532 p.
- Franz, D.R., and Hendler, G.L., 1971, Benthic ecology of a shallow bay: Macrobenthos: Coastal Shallow Water Res. Conf., 2nd Abstr., Baton Rouge; Newark, Del.; Los Angeles, Calif., Oct. 1971, Univ. Press, Univ. Southern Calif., p. 78.
- George, R.Y., and Menzies, R.J., 1973, Deep sea faunal zonation of benthos along Beaufort Bermuda transect in the northwestern Atlantic: Proc. Royal Soc. Edinburg, sect. B, biol., v. 73, 1971-1972, p. 183-194.
- Haedrich, R.L., Rowe, G.T., and Polloni, P.T., 1975, Zonation and faunal composition of epibenthic populations on the continental slope south of New England: Jour. of Marine Research, v. 33, p. 191-212.

Hathaway, J.C., ed., 1966, Data file, Continental Margin Program, Atlantic Coast of the United States; v. 1, Sample collection data: Woods Hole Oceanog. Inst. Ref. no. 66-8, 184 p.

\_\_\_\_\_ 1971, Data file, Continental Margin Program, Atlantic Coast of the United States; v. 2, Sample collection and analytical data: Woods Hole Oceanog. Inst. Ref. no. 71-15, 496 p.

Hollister, C.D., 1973, Atlantic Continental Shelf and Slope of the United States - texture of surface sediments from New Jersey to southern Florida: U.S. Geol. Survey Prof. Paper 529-M, 23 p.

Holme, N.A., and McIntyre, A.D., 1971, Methods for the study of marine benthos: IBP Handbook No. 16, Oxford and Edinburg, Blackwell Sci. Publ., 334 p.

Hülsemann, Jobst, 1967, The continental margin off the Atlantic coast of the United States - Carbonate in sediments, Nova Scotia to Hudson Canyon: Sedimentology, v. 8, no. 2, p. 121-145.

Kaplan, E.H., Welker, J.R., and Kraus, M.G., 1974, Some effects of dredging on populations of macrobenthic organisms: Fishery Bulletin (NOAA), v. 72, n. 2, p. 445-480.

Ketchum, B.H., and Corwin, Nathaniel, 1964, The persistence of "winter" water on continental shelf south of Long Island, New York: Limnology and Oceanography, v. 9, no. 4, p. 467-475.

- Kinner, Peter, Maurer, Don, and Leathem, Wayne, 1974, Benthic invertebrates in Delaware Bay: animal - sediment associations of the dominant species: *Internat. Revue der gesamten Hydrobiologie*, v. 59, no. 5, p. 685-701.
- Lee, R.E., 1944, A quantitative survey of the invertebrate bottom fauna in Menemsha Bight: *Biol. Bull. (Woods Hole)*, v. 86, no. 2, p. 83-97.
- Leidy, Joseph, 1855, Contributions toward a knowledge of the marine invertebrate fauna of the coasts of Rhode Island and New Jersey: *Jour. of the Academy of Natural Sciences of Philadelphia*, v. 3, 2nd ser., reprint, p. 2-19, plates 10, 11.
- Livingstone, Robert Jr., 1965, A preliminary bibliography with KWIC index on the ecology of estuaries and coastal areas of the eastern United States: U.S. Dept. of the Interior, Fish and Wildlife Service, Spec. Sci. Rept. - Fisheries, No. 507, 352 p.
- Maurer, D., Biggs, R., Leathem, W., Kinner, P., Treasure, W., Otley, M., Watling, L., and Klemas, V., 1974, Effect of spoil disposal on benthic communities near the mouth of Delaware Bay: Rept. Del. River Bay Auth., University of Delaware, 200 p.
- McGrath, R.A., 1974, Benthic macrofaunal census of Raritan Bay - Preliminary results: Third Symposium on Hudson River Ecology, The Hudson River Environmental Society, Inc., 40 p.
- Menzies, R.J., Smith, Logan, and Emery, K.O., 1963, A combined underwater camera and bottom grab - A new tool for investigation of deep-sea benthos: *Internat. Revue der gesamten Hydrobiologie*, v. 48, no. 4, 529-545.

Milliman, J.D., 1973, Marine geology in Coastal and offshore environmental inventory, Cape Hatteras to Nantucket Shoals: Univ. of Rhode Island, Marine Publication Series No. 3, Occasional Publ., no. 6, p. 10-1 - 10-91.

National Marine Fisheries Service, 1972, The effects of waste disposal in the New York Bight: NMFS, Middle Atlantic Coastal Fisheries Center, Tech. Rept. 9, 749 p.

National Oceanic and Atmospheric Administration (U.S. Dept. Of Commerce), 1974, Bibliography of the New York Bight, Part 1 - List of citations: Prepared by Environmental Science Information Center, Environmental Data Service, Marine Ecosystems Analysis Program, Office of Coastal Environment, Rockville, Md., 184 p.

\_\_\_\_\_ 1974, Bibliography of the New York Bight, Part 2 - Indexes: Prepared by Environmental Science Information Center, Environmental Data Service, Marine Ecosystems Analysis Program, Office of Coastal Environment, Rockville, Md., 493 p.

O'Connor, J.S., 1972, The benthic macrofauna of Moriches Bay, New York: The Biol. Bull. (Woods Hole), v. 142, no. 1, p. 84-102.

Palmer, H.D., and Lear, D.W., eds., 1973, Environmental survey of an interim ocean dump site, Middle Atlantic Bight: Cruise Report, 1-5 May 1973, EPA 902 - 9 - 001 - A, U.S. Environmental Protection Agency, Region II., Phila., Pa., 134 p.

Parker, R.H., 1974, The study of benthic communities, a model and a review: Elsevier Oceanog. Series, 9, Amsterdam and N.Y., Elsevier Sci. Publ. Co., 260 p.

- Pearce, J.B., 1972, The effects of solid waste disposal on benthic communities in the New York Bight, in Ruivo, Mario, ed., Marine pollution and sealife: Surrey, England, Fishing News (Books) Ltd., p. 404-411.
- \_\_\_\_\_ 1975, The temporal and spatial distribution of benthic macroinvertebrates in the New York Bight: Abstracts, Special Symposium, The Middle Atlantic Continental Shelf and New York Bight, 3-4-5 November 1975, p. 54-56.
- Pearce, J.B., Rogers, Leslie, James, Thomas, Carracciolo, Janice, Halsey, Martha, and McNulty, Knee, 1976, Distribution and abundance of benthic organisms in the outer New York Bight and proposed alternate disposal sites, June 1974 and February 1975: National Oceanic and Atmospheric Administration (NOAA), NOAA Data Rept. ERL MESA - 10, 68 p.
- Phelps, D.K., 1964, Distribution of benthic invertebrates in relationship to the environment of Charlestown Pond. Progress report: Environmental relationships of benthos in salt ponds: Univ. R.I., Grad. School Oceanog., Narragansett Marine Lab., Ref. 64-3, p. 19-54.
- Phillips, F.X., 1972, The ecology of the benthic macro invertebrates of Barnegat Bay, New Jersey: [Abs.] Dissert. Abs. Intl., v. 32, no. 9, p. 5148-B
- Rhoads, D.C., 1963, Rates of sediment reworking by Yoldia limatula in Buzzards Bay, Massachusetts, and Long Island Sound: Jour. Sed. Petrology, v. 33, no. 3, p. 723-727.
- Ross, D.A., 1970, Atlantic Continental Shelf and Slope of the United States - heavy minerals, continental margin, from southern Nova Scotia to northern New Jersey: U.S. Geol Survey Prof. Paper 529-G, 40 p.

Rowe, G.T., 1973, The effects of pollution on the dynamics of the benthos of New York Bight: *Thalassia jugosl*, v. 7, no. 1, p. 353-359.

Rowe, G.T., and Menzel, D.W., 1971, Quantitative benthic samples from the deep Gulf of Mexico with some comments on the measurement of biomass: *Marine Sci. Bull.*, v. 21, p. 556-566.

Rowe, G.T., and Menzies, R.J., 1969, Zonation of large benthic invertebrates in the deep-sea off the Carolinas: *Deep-Sea Research*, v. 16, p. 531-537.

Saila, S.B., (Proj. coord.), 1973, Coastal and offshore environmental inventory: Cape Hatteras to Nantucket Shoals: University of Rhode Island (Kingston, R.I.) Marine Publication Series, no. 2, Chapter bibliographies.

Sanders, H.L., 1956, Oceanography of Long Island Sound 1952-1954. X. The biology of marine bottom communities: *Bingham Oceanog. Collection Bull.*, v. 15, p. 345-414.

\_\_\_\_\_ 1958, Benthic studies in Buzzards Bay. I. Animal-sediment relationships: *Limnology and Oceanography*, v. 3, p. 245-258.

\_\_\_\_\_ 1960, Benthic studies in Buzzards Bay. III. The structure of the soft-bottom community: *Limnology and Oceanography*, v. 5, no. 2, p. 138-153.

Sanders, H.L., and Hessler, R.R., 1969, Ecology of the deep-sea benthos: *Science*, v. 163, p. 1419-1424.

- Sanders, H.L., Hessler, R.R., and Hampson, G.R., 1965, An introduction to the study of deep-sea benthic faunal assemblages along the Gay Head-Bermuda transect: Deep-Sea Research, v. 12, no. 6, p. 845-867.
- Schlee, John, 1973, Atlantic Continental Shelf and Slope of the United States - sediment texture of the northeastern part: U.S. Geol. Survey Prof. Paper 529-L, 64 p.
- Schlee, John, and Pratt, R.M., 1970, Atlantic Continental Shelf and Slope of the United States - gravels of the northeastern part: U.S. Geol. Survey Prof. Paper 529-H, 39 p.
- Smith, W., and McIntyre, A.D., 1954, A spring-loaded bottom sampler: Jour. Mar. Biol. Assoc. U.K., v. 33, p. 257-264.
- Steimle, F.W., and Stone, R.B., 1973, Abundance and distribution of inshore benthic fauna off southwestern Long Island, N.Y.: NOAA Tech. Rept. NMFS SSRF-673, 50 p.
- Stickney, A.P., and Stringer, L.D., 1957, A study of the invertebrate bottom fauna of Greenwich Bay, Rhode Island: Ecology, v. 38, no. 1, p. 111-122.
- Stone, R.B., 1963, A quantitative study of the benthic fauna in lower Chesapeake Bay with emphasis on animal-sediment relationships: M.S. thesis, School of Marine Science, College of William and Mary.
- Stubblefield, W.L., Dicken, Michael, and Swift, D.J.P., 1974, Reconnaissance of bottom sediments on the inner and central New Jersey shelf (MESA Data Report): MESA Rept. No. 1, Marine Ecosystems Analysis Program, Boulder, Colorado, 39 p.
- Sumner, F.B., Osburn, R.C., and Cole, L.J., 1913, A biological survey of the waters of Woods Hole and vicinity: Bull. of the (U.S.) Bur. of Fisheries, v. 31, pt. 1, sect. 1, p. 1-442, 227 charts.

Swift, D.J.P., Duane, D.B., and McKinney, T.F., 1973, Ridge and swale topography of the Middle Atlantic Bight: Secular response to the Holocene hydraulic regime: *Marine Geology*, v. 15, p. 227-247.

Tenore, K.R., 1972, Macrobenthos of the Pamlico River Estuary, North Carolina: *Ecological Monographs*, v. 42, p. 51-69.

Trumbull, J.V.A., 1972, Atlantic Continental Shelf and Slope of the United States - sand size fraction of bottom sediments, New Jersey to Nova Scotia: U.S. Geol. Survey Prof. Paper 529-K, 45 p.

Verill, A.E., 1866, On the polyps and echinoderms of New England, with descriptions of new species: *Proc. Boston Soc. of Nat. Hist.*, v. 10, p. 333-357.

\_\_\_\_\_ 1873, Results of recent dredging expeditions on the coast of New England: *Am. Jour. of Sci. and Arts*, 3d ser., v. 5, p. 98-106.

Walford, L.A., and Wicklund, R.I., 1968, Monthly sea temperature structure from the Florida Keys to Cape Cod: *Am. Geog. Soc., Serial Atlas Marine Environment*, Folio 15.

Watling, L., Leathem, W., Kinner, P., Wethe, C., and Maurer, D., 1974, An evaluation of sewage sludge dumping on the benthos off Delaware Bay: *Marine Pollution Bull.*, v. 5, p. 39-42.

Watling, Les, and Maurer, Don, eds., 1975, Ecological studies on benthic and plankton assemblages in lower Delaware Bay: *Rept. to the Nat'l. Sci. Found. by College of Mar. Studies, Univ. of Delaware*, 630 p.

- Wigley, R.L., and Emery, K.O., 1967, Benthic animals, particularly Hyalinoecia (Annelida) and Ophiomusium (Echinodermata), in sea-bottom photographs from the continental slope: in The Johns Hopkins Oceanographic Studies No. 3, Deep-Sea Photography, John Brackett Hersey, ed., p. 235-249.
- Wigley, R.L., and McIntyre, A.D., 1964, Some quantitative comparisons of offshore meiobenthos and macrobenthos south of Martha's Vineyard: Limnology and Oceanography, v. 9, no. 4, p. 485-493.
- Wigley, R.L., and Stinton, F.C., 1973, Distribution of macroscopic remains of recent animals from marine sediments off Massachusetts: (U.S.) Fishery Bull., v. 71, n. 1, p. 1-40.
- Yentsch, A.E., Carriker, M.R., Parker, R.H., Zullo, V.A., 1966, Marine and estuarine environments, organisms and geology of the Cape Cod region, an indexed bibliography, 1665-1965: Systematics Ecology Program, MBL (Marine Biological Laboratory), Woods Hole, Mass., 178 p.

APPENDIX

Appendix Table A-1.--Number of samples within each depth range class  
in each subarea and for the entire Middle Atlantic  
Bight Region.

Depth range	Subarea			Entire region
	Southern New England	New York Bight	Chesapeake Bight	
<u>m</u>	<u>No.</u>	<u>No.</u>	<u>No.</u>	<u>No.</u>
0 - 24	35	46	84	165
25 - 49	27	48	48	123
50 - 99	56	47	15	118
100 - 199	19	9	6	34
200 - 499	14	8	6	28
500 - 999	8	7	10	25
1,000 - 1,999	11	10	13	24
2,000 - 3,080	16	12	8	36
Total	186	187	190	563

Appendix Table A-2.--Number of samples for each bottom sediment type in each subarea and for the entire Middle Atlantic Bight Region.

Bottom sediments	Subarea			Entire region
	Southern New England	New York Bight	Chesapeake Bight	
	<u>No.</u>	<u>No.</u>	<u>No.</u>	<u>No.</u>
Gravel	3	0	0	3
Sand-gravel	11	5	2	18
Shell	1	0	3	4
Sand-shell	1	16	27	44
Sand	83	118	84	285
Silty sand	52	18	24	94
Silt	25	16	28	69
Clay	10	14	22	46
Total	186	187	190	563

Appendix Table A-3.--Number of samples for each class of sediment organic carbon in each subarea and for the entire Middle Atlantic Bight Region.

Organic carbon	Subarea			Entire region
	Southern New England	New York Bight	Chesapeake Bight	
<u>%</u>	<u>No.</u>	<u>No.</u>	<u>No.</u>	<u>No.</u>
0.01 - 0.4	93	139	117	349
0.5 - 0.9	55	29	26	110
1.0 - 1.4	14	9	17	40
1.5 - 1.9	4	6	15	25
2.0 - 2.9	1	4	4	9
3.0 - 4.9	0	0	9	9
5.0 - 7.2	0	0	1	1
No data	19	0	1	20
Total	186	187	190	563

Appendix Table A-4.--Number of samples within each water temperature range class in each subarea and for the entire Middle Atlantic Bight Region.

Temperature range	Subarea			Entire region
	Southern New England	New York Bight	Chesapeake Bight	
<u>°C</u>	<u>No.</u>	<u>No.</u>	<u>No.</u>	<u>No.</u>
0 - 3.9	46	36	28	110
4.0 - 7.9	7	5	5	17
8.0 - 11.9	12	16	5	33
12.0 - 15.9	52	42	8	102
16.0 - 19.9	31	32	16	79
20.0 - 23.9	28	52	74	154
24.0 +	10	4	54	68
Total	186	187	190	563