


SPECIES: BLACK SEA BASS



SPECIES: FLUKE





Figure 1B. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.
 Top: black sea bass (*Centropristis striata*); bottom: fluke (*Paralichthys dentatus*).

SPECIES: SCUP



SPECIES: AMERICAN PLAICE





Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: scup (*Stenotomus chrysops*); bottom: American plaice (*Hippoglossoides platessoides*).

SPECIES: ATLANTIC COD



SPECIES: ATLANTIC HALIBUT





Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: Atlantic cod (*Gadus morhua*); bottom: Atlantic halibut (*Hippoglossus hippoglossus*).

SPECIES: ATLANTIC WOLFFISH



SPECIES: HADDOCK





Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: Atlantic wolffish (*Anarhichas lupus*); bottom: haddock (*Melanogrammus aeglefinus*).

SPECIES: OCEAN POUT



SPECIES: POLLOCK





Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: ocean pout (*Zoarces americanus*); bottom: pollock (*Pollachius virens*).

SPECIES: REDFISH



SPECIES: WHITE HAKE





Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: redfish (*Sebastes fasciatus*); bottom: white hake (*Urophycis tenuis*).

SPECIES: WINDOWPANE FLOUNDER



SPECIES: WINTER FLOUNDER





Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: windowpane flounder (*Scophthalmus aquosus*); bottom: winter flounder (*Pseudopleuronectes americanus*).

SPECIES: WITCH FLOUNDER



SPECIES: YELLOWTAIL FLOUNDER





Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: witch flounder (*Glyptocephalus cynoglossus*); bottom: yellowtail flounder (*Limanda ferruginea*).

SPECIES: OFFSHORE HAKE



SPECIES: RED HAKE





Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: offshore hake (*Merluccius albidus*); bottom: red hake (*Urophycis chuss*).

SPECIES: SILVER HAKE



SPECIES: ATLANTIC MACKEREL





Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: silver hake (*Merluccius bilinearis*); bottom: Atlantic mackerel (*Scomber scombrus*).

SPECIES: BUTTERFISH



SPECIES: NORTHERN SHORTFIN SQUID




Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: butterfish (*Peprilus triacanthus*); bottom: northern shortfin squid (*Illex illecebrosus*).

SPECIES: LONGFIN INSHORE SQUID




Figure 1B, continued. Percentage of Vessel Trip Report landings (kept) and estimated discards (Total, left pie) and the percentage of estimated discards by fleet (Discards, right pie) for the 23 individual species that compose the 14 species groups, based on July 2015 through June 2016 data. Because percentages have been rounded, they may not always sum to 100%. See Appendix Table 4 for fleet abbreviations.

Top: Longfin inshore squid (*Doryteuthis [Amerigo] pealeii*).