


## SHORT-FINNED PILOT WHALE (*Globicephala macrorhynchus*): Western North Atlantic Stock

### STOCK DEFINITION AND GEOGRAPHIC RANGE

There are two species of pilot whales in the western North Atlantic - the long-finned pilot whale, *Globicephala melas melas*, and the short-finned pilot whale, *G. macrorhynchus*. These species are difficult to differentiate at sea and cannot be reliably visually identified during either abundance surveys or observations of fishery mortality without high-quality photographs (Rone and Pace 2012); therefore, the ability to separately assess the two species in U.S. Atlantic waters is complex and requires additional information on seasonal spatial distribution. Pilot whales (*Globicephala* sp.) in the western North Atlantic occur primarily along the continental shelf break from Florida to the Nova Scotia Shelf (Mullin and Fulling 2003). Long-finned and short-finned pilot whales overlap spatially along the mid-Atlantic shelf break between Delaware and the southern flank of Georges Bank (Payne and Heinemann 1993; Rone and Pace 2012). Long-finned pilot whales have occasionally been observed stranded as far south as South Carolina, and short-finned pilot whales have occasionally been observed stranded as far north as Massachusetts (Pugliares *et al.* 2016). The exact latitudinal ranges of the two species remain uncertain. However, south of Cape Hatteras most pilot whale sightings are expected to be short-finned pilot whales, while north of ~42°N most pilot whale sightings are expected to be long-finned pilot whales (Figure 1; Garrison and Rosel 2017). Short-finned pilot whales are also documented along the continental shelf and continental slope in the northern Gulf of Mexico (Hansen *et al.* 1996;



**Figure 1.** Distribution of long-finned (open symbols), short-finned (black symbols), and possibly mixed (gray symbols; could be either species) pilot whale sightings from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1998, 1999, 2002, 2004, 2006, 2007, 2011, and 2016. The inferred distribution of the two species is preliminary and is valid for June-August only. Isobaths are the 1,000-m and 3,000-m depth contours. The U.S. EEZ is also displayed in green.

Mullin and Hoggard 2000; Mullin and Fulling 2003), and are known from the wider Caribbean (Bernard and Riley 1999). Five short-finned pilot whales tagged in the Great Bahama Canyon, northern Bahamas, were tracked into the Gulf Stream and moved north to deep waters off the coast of central and northern Florida (Claridge *et al.* 2015), suggesting the potential for connectivity between pilot whales in the southern U.S. range of this stock and the Caribbean. However, none of the tagged whales moved north of South Carolina (Claridge *et al.* 2015) which could suggest multiple populations in the stock range (e.g., a northern and a southern population), or simply that tag duration was too short to detect broader movements. Two tagged and released individuals from a May 2011 mass stranding of 23 short-finned pilot whales in the Florida Keys travelled to waters off South Carolina, and one subsequently moved to waters between Cuba and Haiti (Wells *et al.* 2013). Short-finned pilot whales tagged during a 1977 mass stranding near Jacksonville were recovered off South Carolina (Irvine *et al.* 1979). It is not known how representative of normal species patterns any of these movements are. An analysis of stock structure within the western North Atlantic Stock has not been completed so there are insufficient data to determine whether there are multiple demographically-independent populations within this stock. Continued studies to evaluate genetic population structure in short-finned pilot whales throughout the region will improve understanding of stock structure. Pending these results, the *Globicephala macrorhynchus* population occupying U.S. Atlantic waters is considered separate from both the northern Gulf of Mexico stock and short-finned pilot whales occupying Caribbean waters.

### POPULATION SIZE

The best available estimate for short-finned pilot whales in the western North Atlantic is 28,924 (CV=0.24; Table 1; Palka 2012; Garrison 2016; Garrison and Rosel 2017; Garrison and Palka 2018). This estimate is from summer 2016 surveys covering waters from central Florida to Georges Bank. Pilot whale sightings from vessel surveys were strongly concentrated along the continental shelf break; however, pilot whales were also observed over the continental slope in waters associated with the Gulf Stream (Figure 1). The best available abundance estimates are from shipboard surveys conducted during the summer of 2016 because these are the most recent surveys covering the full range of short-finned pilot whales in U.S. Atlantic waters. Because long-finned and short-finned pilot whales are difficult to distinguish at sea, sightings data are reported as *Globicephala sp.* These survey data have been combined with an analysis of the spatial distribution of the two pilot whale species based on genetic analyses of biopsy samples to derive separate abundance estimates for each species (Garrison and Rosel 2017).

### Earlier Estimates

Please see Appendix IV for a summary of abundance estimates including earlier estimates and survey descriptions. Due to changes in survey methodology, these historical data should not be used to make comparisons with more current estimates. In addition, as recommended in the GAMMS II Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable for the determination of a current PBR.

### Recent surveys and abundance estimates for *Globicephala sp.*

For waters between central Virginia and the lower Bay of Fundy, an abundance estimate of 11,865 (CV=0.57) *Globicephala sp.* was generated from aerial and shipboard surveys conducted during June–August 2011 (Palka 2012). The aerial portion covered 6,850 km of trackline over waters north of New Jersey between the coastline and the 100-m depth contour through the U.S. and Canadian Gulf of Maine, and up to and including the lower Bay of Fundy. Pilot whales were not observed during the aerial portion of the survey. The shipboard portion covered 3,811 km of trackline between central Virginia and Massachusetts in waters deeper than the 100-m depth contour out to beyond the U.S. Exclusive Economic Zone (EEZ). Estimation of abundance was based on the independent observer approach, which allows estimation of abundance corrected for perception bias of the detected species, assuming point independence (Laake and Borchers 2004) and calculated using the mark-recapture distance sampling option in the computer program Distance (version 6.0, release 2, Thomas *et al.* 2009). The vessel portion of this survey included habitats where both short-finned and long-finned pilot whales occur. Short-finned pilot whales are not predicted to occur north of Georges Bank. A logistic regression (see next section) was used to estimate the abundance of short-finned pilot whales from this survey as 4,569 (CV=0.57).

For waters between central Virginia and central Florida, an abundance estimate of 16,946 (CV=0.43) *Globicephala sp.* was generated from a shipboard survey conducted during June–August 2011 (Garrison 2016). This shipboard survey included shelf-break and inner continental slope waters deeper than the 50-m depth contour within the U.S. EEZ. The survey employed two independent visual teams searching with 25x150 “bigeye” binoculars. A total of 4,445 km of trackline was surveyed. The majority of pilot whale sightings occurred along the continental shelf break north of Cape Hatteras, North Carolina, with a lower number of sightings over the continental slope in

the southern portion of the survey. Estimation of pilot whale abundance was based on the independent observer approach, which allows estimation of abundance corrected for perception bias of the detected species, assuming point independence (Laake and Borchers 2004) and calculated using the mark-recapture distance sampling option in the computer program Distance (version 6.0, release 2, Thomas *et al.* 2009). A logistic regression (see next section) was used to estimate the abundance of short-finned pilot whales from this survey. The regression indicated this survey included habitats expected to exclusively contain short-finned pilot whales resulting in an abundance estimate of 16,946 (CV=0.43) short-finned pilot whales from this survey.

Abundance estimates of 8,166 (CV=0.31) and 25,114 (CV=0.27) *Globicephala* sp. were generated from vessel surveys conducted in the northeast and southeast U.S., respectively, during the summer of 2016. The Northeast survey was conducted during 27 June – 25 August and consisted of 5,354 km of on-effort trackline. The majority of the survey was conducted in waters north of 38°N latitude and included tracklines along the shelf break and offshore to the U.S. EEZ. Pilot whale sightings were concentrated along the shelf-break between the 1,000-m and 2,000-m isobaths and along Georges Bank (NMFS 2017). The Southeast vessel survey covered waters from Central Florida to approximately 38°N latitude between the 100-m isobaths and the U.S. EEZ during 30 June – 19 August. A total of 4,399 km of trackline was covered on effort. Pilot whales were observed in high densities along the shelf-break between Cape Hatteras and New Jersey and also in waters further offshore in the mid-Atlantic and off the coast of Florida (NMFS 2017; Garrison and Palka 2018). Both the Northeast and Southeast surveys utilized two visual teams and an independent observer approach to estimate detection probability on the trackline (Laake and Borchers 2004). Mark-recapture distance sampling was used to estimate abundance. A logistic regression model (see next section) was used to estimate the abundance of short-finned pilot whales from these surveys. For the northeast survey, this resulted in an abundance estimate of 3,810 (CV=0.42) short-finned pilot whales. In the southeast, the model indicated that this survey included habitats expected to exclusively contain short-finned pilot whales resulting in an abundance estimate of 25,114 (CV=0.27).

#### **Spatial Distribution and Abundance Estimates for *Globicephala macrorhynchus***

Pilot whale biopsy samples were collected during summer months (June–August) from South Carolina to the southern flank of Georges Bank between 1998 and 2007. These samples were identified to species using phylogenetic analysis of mitochondrial DNA sequences. Samples from stranded specimens that were morphologically identified to species were used to assign clades in the phylogeny to species and thereby identify all survey samples. The probability of a sample being from a short-finned (or long-finned) pilot whale was evaluated as a function of sea surface temperature, latitude, and month using a logistic regression. This analysis indicated that the probability of a sample coming from a short-finned pilot whale was near zero at water temperatures <22°C, and near one at temperatures >25°C. The probability of being a short-finned pilot whale also decreased with increasing latitude. Spatially, during summer months, this regression model predicted that all pilot whales observed in offshore waters near the Gulf Stream are most likely short-finned pilot whales. The area of overlap between the two species occurs primarily along the shelf break between 38°N and 40°N latitude (Garrison and Rosel 2017). This model was used to partition the abundance estimates from surveys conducted during the summers of 2011 and 2016. The sightings from the shipboard surveys covering waters from Florida to New Jersey were predicted to consist entirely of short-finned pilot whales. The vessel portion of the northeast surveys from New Jersey to the southern flank of Georges Bank included waters along the shelf break and waters further offshore extending to the U.S. EEZ. Pilot whales were observed in both areas during the survey. Along the shelf break, the model predicted a mixture of both species, but the sightings in offshore waters near the Gulf Stream were again predicted to consist predominantly of short-finned pilot whales (Garrison and Rosel 2017). The best abundance estimate for short-finned pilot whales is thus the sum of the southeast survey estimate (25,114; CV=0.27) and the estimated number of short-finned pilot whales from the northeast vessel survey (3,810; CV=0.42). The best available abundance estimate is thus 28,924 (CV=0.24).

**Table 1. Summary of recent abundance estimates for the western North Atlantic short-finned pilot whale (*Globicephala macrorhynchus*) by month, year, and area covered during each abundance survey, and resulting abundance estimate ( $N_{best}$ ) and coefficient of variation (CV).**

Month/Year	Area	$N_{best}$	CV
Jun–Aug 2011	central Virginia to Georges Bank	4,569	0.57
Jun–Aug 2011	central Florida to central Virginia	16,946	0.43

Month/Year	Area	N <sub>best</sub>	CV
Jun–Aug 2011	central Florida to Georges Bank (COMBINED)	21,515	0.37
Jun–Aug 2016	New Jersey to Georges Bank	3,810	0.42
Jun–Aug 2016	central Florida to New Jersey	25,114	0.27
Jun–Aug 2016	central Florida to Georges Bank (COMBINED)	28,924	0.24

### Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for western North Atlantic *Globicephala macrorhynchus* is 28,924 animals (CV=0.24). The minimum population estimate is 23,637.

### Current Population Trend

There are three available coastwide abundance estimates for short-finned pilot whales from the summers of 2004, 2011, and 2016. Each of these is derived from vessel surveys with similar survey designs and all three used the two-team independent observer approach to estimate abundance. The southeast component of these surveys all were expected to contain exclusively short-finned pilot whales, and the logistic regression model was used to partition pilot whale sightings from the northeast portion of the survey between the short-finned and long-finned species based upon habitat characteristics. The resulting estimates were 24,674 (CV=0.52) in 2004, 21,515 (CV=0.36) in 2011, and 28,924 (CV=0.24) in 2016 (Garrison and Palka 2018). A generalized linear model indicated no significant trend in these abundance estimates. The key uncertainty is the assumption that the logistic regression model accurately represents the relative distribution of short-finned vs. long-finned pilot whales in each year.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

### POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size for short-finned pilot whales is 23,637. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor is 0.5 because the stock’s status relative to optimum sustainable population (OSP) is unknown and the CV of the average mortality estimate is less than 0.3 (Wade and Angliss 1997). PBR for the western North Atlantic short-finned pilot whale is 236.

### ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

The total annual human-caused mortality and serious injury for this stock during 2012–2016 is unknown. The estimated mean annual fishery-related mortality and serious injury during 2012–2016 due to the pelagic longline fishery was 168 short-finned pilot whales (CV=0.13; Table 2). Uncertainty in this estimate arises because it incorporates a logistic regression model to predict the species of origin (long-finned or short-finned pilot whale) for each bycaught whale. The statistical uncertainty in the assignment to species is incorporated into the abundance estimates; however, the analysis assumes that the collected biopsy samples adequately represent the distribution of the two species and that the resulting model correctly predicts shifts in distribution in response to changes in environmental conditions. In addition to observed takes in the pelagic longline fishery, there was a self-reported take in 2013 in the unobserved hook and line fishery. This unobserved take renders the estimate of total annual fishery-caused mortality and serious injury an underestimate.

In bottom trawl, mid-water trawl, and gillnet fisheries, pilot whale mortalities were observed north of 40°N latitude in areas expected to have only long-finned pilot whales. Takes and bycatch estimates for these fisheries are therefore attributed to the long-finned pilot whale stock.

### **Fishery Information**

There are four commercial fisheries that interact, or that potentially could interact, with this stock in the Atlantic Ocean. These include two Category I fisheries (Atlantic Ocean, Caribbean, Gulf of Mexico large pelagics longline and Atlantic Highly Migratory Species longline fisheries) and two Category III fisheries (U.S. Atlantic tuna purse seine and Atlantic Ocean, Gulf of Mexico, Caribbean commercial passenger fishing vessel (hook and line) fisheries). All recent gillnet and trawl interactions have been assigned to long-finned pilot whales using model-based predictions. Detailed fishery information is reported in Appendix III.

### **Earlier Interactions**

See Appendix V for information on historical takes.

### **Longline**

The Atlantic Ocean, Caribbean, Gulf of Mexico large pelagics longline fishery operates in the U.S. Atlantic (including Caribbean) and Gulf of Mexico EEZ, and pelagic swordfish, tunas and billfish are the target species. The estimated annual average serious injury and mortality attributable to the Atlantic Ocean large pelagics longline fishery for the five-year period from 2012 to 2016 was 168 short-finned pilot whales (CV=0.13; Table 2). During 2012–2016, 92 serious injuries were observed in the following fishing areas of the North Atlantic: Florida East Coast, Mid-Atlantic Bight, Northeast Coastal, and South Atlantic Bight. During 2012–2016, one mortality was observed (in 2016) in the Florida East Coast fishing area (Garrison and Stokes 2013; 2014; 2016; 2017; 2019).

Prior to 2014, estimated bycatch in the pelagic longline fishery was assigned to the short-finned pilot whale stock because the observed interactions all occurred at times and locations where available data indicated that long-finned pilot whales were very unlikely to occur. Specifically, the highest bycatch rates of undifferentiated pilot whales were observed during September–November along the mid-Atlantic coast (south of 38°N; Garrison 2007), and biopsy data collected in this area during October–November 2011 indicated that only short-finned pilot whales occurred in this region (Garrison and Rosel 2017). Similarly, all genetic data collected from interactions in the pelagic longline fishery have indicated interactions with short-finned pilot whales. However, during 2014–2016, pilot whale interactions (including serious injuries) were observed further north and along the southern flank of Georges Bank. Therefore, the logistic regression model (described above in 'Spatial Distribution and Abundance Estimates for *Globicephala macrorhynchus*') was applied to estimate the probability that these interactions were from short-finned vs. long-finned pilot whales (Garrison and Rosel 2017). Due to high water temperatures (ranging from 22 to 25°C) at the time of the observed takes, these interactions were estimated to have a >90% probability of coming from short-finned pilot whales. The estimated probability was used to apportion the estimated serious injury and mortality from 2014 to 2016 in the pelagic longline fishery between the short-finned and long-finned pilot whale stocks (Garrison and Stokes 2016; 2017; 2019).

Between 1992 and 2004, most of the marine mammal bycatch in the U.S. pelagic longline fishery was recorded in U.S. Atlantic EEZ waters between South Carolina and Cape Cod (Garrison 2007). From January to March, observed bycatch was concentrated on the continental shelf edge northeast of Cape Hatteras, North Carolina. During April–June, bycatch was recorded in this area as well as north of Hydrographer Canyon in water over 1,000 fathoms (1830 m) deep. During the July–September period, observed takes occurred on the continental shelf edge east of Cape Charles, Virginia, and on Block Canyon slope in over 1,000 fathoms of water. October–December bycatch occurred between the 20- and 50-fathom (37- and 92-m) isobaths between Barnegat Bay, New Jersey, and Cape Hatteras, North Carolina.

The Atlantic Highly Migratory Species longline fishery operates outside the U.S. EEZ. No takes of short-finned pilot whales within high seas waters of the Atlantic Ocean have been observed or reported thus far.

See Table 2 for bycatch estimates and observed mortality and serious injury for the current five-year period, and Appendix V for historical estimates of annual mortality and serious injury.

**Table 2. Summary of the incidental mortality and serious injury of short-finned pilot whales (*Globicephala macrorhynchus*) by the pelagic longline commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the annual observed serious injury and mortality recorded by on-board observers, the annual estimated serious injury and mortality, the combined annual estimates of serious injury and mortality (Estimated Combined Mortality), the estimated CV of the combined annual mortality estimates (Est. CVs) and the mean of the combined mortality estimates (CV in parentheses).**

Fishery	Years	Vessels <sup>a</sup>	Data Type <sup>b</sup>	Percent Observer Coverage <sup>c</sup>	Observed Serious Injury	Observed Mortality	Est. Serious Injury	Est. Mortality	Est. Combined Mortality	Est. CVs	Mean Annual Mortality
Pelagic Longline	2012	82	Obs. Data, Logbook	7	14	0	170	0	170	0.33	168 (0.13)
	2013	79		9	13	0	124	0	124	0.32	
	2104	78		10	19	0	233	0	233	0.24	
	2015	74		12	32	0	200	0	200	0.24	
	2016	60		15	14	1	106	5.1	111	0.31	

<sup>a</sup> Number of vessels in the fishery is based on vessels reporting effort to the pelagic longline logbook.

<sup>b</sup> Observer data (Obs. Data) are used to measure bycatch rates and the data are collected within the Northeast Fisheries Observer Program (NEFOP) and the Southeast Pelagic Longline Observer Program.

<sup>c</sup> Percentage of sets observed

### Hook and Line

During 2012–2016, there was one self-reported take (in 2013) in which a short-finned pilot whale was hooked and entangled by a charterboat fisherman. The animal was released alive but considered seriously injured (Maze-Foley and Garrison 2016).

### Other Mortality

Pilot whales have a propensity to mass strand throughout their range, but the role of human activity in these events is unknown. Between two and 168 pilot whales have stranded annually, either individually or in groups, along the eastern U.S. seaboard since 1980 (NMFS 1993; stranding databases maintained by NMFS NER, NEFSC and SEFSC). During 2012–2016, 39 short-finned pilot whales (*Globicephala macrorhynchus*) and one pilot whale not specified to the species level (*Globicephala* sp.) were reported stranded between Massachusetts and Florida (Table 3; Northeast Regional Marine Mammal Stranding Network; Southeast Regional Marine Mammal Stranding Network; NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 28 April June 2017 (SER) and 5 May 2017 (NER)).

**Table 3. Short-finned pilot whale (*Globicephala macrorhynchus* [SF] and *Globicephala* sp. [Sp]) strandings along the Atlantic coast, 2012–2016. Strandings which were not reported to species have been reported as *Globicephala* sp. The level of technical expertise among stranding network personnel varies, and given the potential difficulty in correctly identifying stranded pilot whales to species, reports to specific species should be viewed with caution. Data are from the NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 28 April June 2017 (SER) and 5 May 2017 (NER).**

STATE	2012-SF	2012-Sp	2013-SF	2013-Sp	2014-SF	2014-Sp	2015-SF	2015-Sp	2016-SF	2016-Sp	TOT AL-SF	TOT AL-Sp
Massachusetts	0	0	0	0	0	0	0	0	1	0	1	0
North Carolina	1 <sup>a</sup>	0	0	0	3	0	2	0	0	0	6	0
South Carolina	3 <sup>b</sup>	1	1	0	2	0	0	0	0	0	6	1
Georgia	0	0	0	0	0	0	1	0	0	0	1	0
Florida	23 <sup>c</sup>	0	0	0	0	0	2	0	0	0	25	0
<b>TOTALS</b>	<b>27</b>	<b>1</b>	<b>1</b>	<b>0</b>	<b>5</b>	<b>0</b>	<b>5</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>39</b>	<b>1</b>

<sup>a</sup> Signs of fishery interaction were observed for this short-finned pilot whale stranding.

<sup>b</sup> Signs of fishery interaction were observed for 2 of these short-finned pilot whale strandings.

<sup>c</sup> These animals mass stranded alive in September 2012.

One short-finned pilot whale stranding was reported as far north as Cape Cod, Massachusetts (2016); the remaining strandings occurred from North Carolina southward (Table 3).

During 2012–2016, several fishery interactions were documented in stranded pilot whales along the U.S. Atlantic coast. In 2012, three short-finned pilot whales had evidence of fishery interactions, two of them in South Carolina and one in North Carolina. During 2012–2016, no evidence of other human interactions was documented for stranded pilot whales. These strandings are not included in the estimate of total human-caused mortality and serious injury.

Stranding data probably underestimate the extent of human and fishery-related mortality and serious injury, particularly for offshore species such as pilot whales, because not all of the whales that die or are seriously injured in human interactions wash ashore, or, if they do, they are not all recovered (Peltier *et al.* 2012; Wells *et al.* 2015). Additionally, not all carcasses will show evidence of human interaction, entanglement or other fishery-related interaction due to decomposition, scavenger damage, etc. (Byrd *et al.* 2014). Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of human interaction.

### HABITAT ISSUES

The chronic impacts of contaminants (polychlorinated biphenyls [PCBs] and chlorinated pesticides [DDT, DDE, dieldrin, etc.]) on marine mammal reproduction and health are of concern (e.g., Schwacke *et al.* 2002; Jepson *et al.* 2016; Hall *et al.* 2018). Moderate levels of these contaminants have been found in pilot whale blubber (Taruski *et al.* 1975; Muir *et al.* 1988; Weisbrod *et al.* 2000). Weisbrod *et al.* (2000) examined polychlorinated biphenyl and chlorinated pesticide concentrations in bycaught and stranded pilot whales in the western North Atlantic. Contaminant levels were similar to or lower than levels found in other toothed whales in the western North Atlantic, perhaps because they are feeding further offshore than other species (Weisbrod *et al.* 2000). Dam and Bloch (2000) found very high PCB levels in long-finned pilot whales in the Faroes. Also, high levels of toxic metals (mercury, lead, cadmium) and selenium were measured in pilot whales harvested in the Faroe Island drive fishery (Nielsen *et al.* 2000). However, the population effect of the observed levels of such contaminants on this stock is unknown.

### STATUS OF STOCK

The short-finned pilot whale is not listed as threatened or endangered under the Endangered Species Act, and the western North Atlantic stock is not a strategic stock under the MMPA because the mean annual human-caused mortality and serious injury does not exceed PBR. The status of this stock relative to OSP in the U.S. Atlantic EEZ is unknown. Total U.S. fishery-related mortality and serious injury attributed to short-finned pilot whales exceeds 10% of the calculated PBR and therefore cannot be considered to be insignificant and approaching zero mortality and serious injury rate. There is no evidence for a trend in population size for this stock. Should there be multiple demographically-independent stocks within this stock's range, the geographically-concentrated nature of the fishery-related mortality and serious injury could mean that the mortality is impacting one stock more than the other.

### REFERENCES CITED

- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade. 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp. Available from: <http://www.nmfs.noaa.gov/pr/pdfs/sars/guidelines1995.pdf>
- Bernard, H. J. and S.B. Reilly. 1999. Pilot whales *Globicephala* Lesson, 1828. Pages 245–279 in: S. H. Ridgway and R. Harrison (eds), Handbook of marine mammals, Vol. 6: The second book of dolphins and the porpoises. Academic Press, San Diego, CA, USA.
- Byrd, B.L., A.A. Hohn, G.N. Lovewell, K.M. Altman, S.G. Barco, A. Friedlaender, C.A. Harms, W.A. McLellan, K.T. Moore, P.E. Rosel and V.G. Thayer. 2014. Strandings illustrate marine mammal biodiversity and human impacts off the coast of North Carolina, USA. Fish. Bull. 112:1–23.
- Claridge, D., G. M. Ylitalo, D. Herman, J. Durban and K. Parsons. 2015. Behavioral ecology of deep diving odontocetes in the Bahamas. Final Report SERDP Project Number: RC-2114. 114 pp. Available from: [https://swfsc.noaa.gov/uploadedFiles/Events/Meetings/MMT\\_2015/Documents/4.4%20Ppr%202015\\_SERDP%20Draft%20Final%20Report\\_RC2114.pdf](https://swfsc.noaa.gov/uploadedFiles/Events/Meetings/MMT_2015/Documents/4.4%20Ppr%202015_SERDP%20Draft%20Final%20Report_RC2114.pdf)
- Dam, M. and D. Bloch. 2000. Screening of mercury and persistent organochlorine pollutants in long-finned pilot whale (*Globicephala melas*) in the Faroe Islands. Mar. Poll. Bull. 40(12):1090–1099.
- Garrison, L.P. 2007.

- Interactions between marine mammals and longline fishing gear in the U.S. Atlantic Ocean between 1992 and 2004. *Fish. Bull.* 105(3):408–417.
- Garrison, L.P. 2016. Abundance of marine mammals in waters of the U.S. East Coast during summer 2011. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, FL 33140. PRBD Contribution # PRBD-2016-08, 21 pp.
- Garrison, L.P. and D. Palka. 2018. Abundance of short-finned pilot whales along the U.S. east coast from summer 2016 vessel surveys. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, FL 33140. PRBD Contribution # PRBD-2018-07, 17 pp.
- Garrison, L.P. and P.E. Rosel. 2017. Partitioning short-finned and long-finned pilot whale bycatch estimates using habitat and genetic information. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, FL 33140. PRBD Contribution # PRBD-2016-17, 24 pp.
- Garrison, L.P. and L. Stokes. 2013. Estimated bycatch of marine mammals and sea turtles in the U.S. Atlantic pelagic longline fleet during 2012. NOAA Tech. Memo. NMFS-SEFSC-655. 62 pp.
- Garrison, L.P. and L. Stokes. 2014. Estimated bycatch of marine mammals and sea turtles in the U.S. Atlantic pelagic longline fleet during 2013. NOAA Tech. Memo. NMFS-SEFSC-667. 61 pp.
- Garrison, L.P. and L. Stokes. 2016. Estimated bycatch of marine mammals and sea turtles in the U.S. Atlantic pelagic longline fleet during 2014. NOAA Tech. Memo. NMFS-SEFSC-696, 62 pp.
- Garrison, L.P. and L. Stokes. 2017. Estimated bycatch of marine mammals and sea turtles in the U.S. Atlantic pelagic longline fleet during 2015. NOAA Tech. Memo. NMFS-SEFSC-709, 61 pp.
- Garrison, L.P. and L. Stokes. 2019. Estimated bycatch of marine mammals and sea turtles in the U.S. Atlantic pelagic longline fleet during 2016. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, FL 33140. PRBD Contribution # PRBD-2019-01. 62 pp.
- Hall, A.J., B.J. McConnell, L.J. Schwacke, G.M. Ylitalo, R. Williams and T.K. Rowles. 2018. Predicting the effects of polychlorinated biphenyls on cetacean populations through impacts on immunity and calf survival. *Environ. Poll.* 233:407–418.
- Hansen, L.J., K.D. Mullin, T.A. Jefferson and G.P. Scott. 1996. Visual surveys aboard ships and aircraft. Pages 55–132 *in*: R.W. Davis and G.S. Fargion (eds.) Distribution and abundance of marine mammals in the north-central and western Gulf of Mexico: Final report. Volume II: Technical report. OCS Study MMS 96- 0027. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA.
- Jepson, P.D., R. Deaville, J.L. Barber, A. Aguilar, A. Borrell, S. Murphy, J. Barry, A. Brownlow, J. Barnett, S. Berrow and A.A. Cunningham. 2016. PCB pollution continues to impact populations of orcas and other dolphins in European waters. *Sci. Rep.-U.K.* 6:18573.
- Laake, J.L. and D.L. Borchers. 2004. Methods for incomplete detection at distance zero. Pages 108–189 *in*: S.T. Buckland, D.R. Andersen, K.P. Burnham, J.L. Laake, and L. Thomas, (eds.) Advanced distance sampling. Oxford University Press, New York.
- Maze-Foley, K. and L.P. Garrison. 2016. Serious injury determinations for small cetaceans off the southeast U.S. coast, 2013. Southeast Fisheries Science Center, Protected Resources and Biodiversity Division, 75 Virginia Beach Dr., Miami, FL 33140. PRBD Contribution # PRBD-2016-06, 12 pp.
- Muir, D.C.G., R. Wagemann, N.P. Grift, R.J. Norstrom, M. Simon and J. Lien. 1988. Organochlorine chemical and heavy metal contaminants in white-beaked dolphins (*Lagenorhynchus albirostris*) and pilot whales (*Globicephala melaena*) from the coast of Newfoundland, Canada. *Arch. Environ. Contam. Toxicol.* 17(5): 613–629.
- Mullin, K.D. and G.L. Fulling. 2003. Abundance of cetaceans in the southern U.S. North Atlantic Ocean during summer 1998. *Fish. Bull.* 101(3):603–613.
- Mullin, K.D. and W. Hoggard. 2000. Visual surveys of cetaceans and sea turtles from aircraft and ships. Pages 111–172 *in*: R.W. Davis, W.E. Evans and B. Würsig (eds.) Cetaceans, sea turtles and seabirds in the northern Gulf of Mexico: Distribution, abundance and habitat associations. Volume II: Technical report. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans. OCS Study MMS 96-0027.
- Nielsen, J.B., F. Nielsen, P.-J. Jorgensen and P. Grandjean. 2000. Toxic metals and selenium in blood from pilot whales (*Globicephala melas*) and sperm whales (*Physeter catodon*). *Mar. Poll. Bull.* 40(4):348–35.
- NMFS. 1993. Status of fishery resources off the northeastern United States for 1993. NOAA Tech. Memo. NMFS-F/NEC-101, 140 pp. Available from: <http://www.nefsc.noaa.gov/publications/tm/pdfs/tmfneec101.pdf>

- NMFS. 2017. Annual report of a comprehensive assessment of marine mammal, marine turtle, and seabird abundance and spatial distribution in US waters of the western North Atlantic Ocean – AMAPPS II. Available from: <https://www.nefsc.noaa.gov/psb/AMAPPS/>
- Palka, D. 2012. Cetacean abundance estimates in US northwestern Atlantic Ocean waters from summer 2011 line transect survey. U.S. Dept. Commer., Northeast Fish. Sci. Cent. Ref. Doc. 12-29, 37 pp. Available from: <http://www.nefsc.noaa.gov/nefsc/publications/>
- Payne, P.M. and D.W. Heinemann. 1993. The distribution of pilot whales (*Globicephala* sp.) in shelf/shelf edge and slope waters of the northeastern United States, 1978-1988. Rep. Int. Whal. Comm. (Special Issue) 14:51–68.
- Peltier, H., W. Dabin, P. Daniel, O. Van Canneyt, G. Dorémus, M. Huon and V. Ridoux. 2012. The significance of stranding data as indicators of cetacean populations at sea: modelling the drift of cetacean carcasses. Ecol. Indic. 18:278–290.
- Pugliares, K.R., T.W. French, G.S. Jones, M.E. Niemeyer, L.A. Wilcox and B.J. Freeman. 2016. First records of the short-finned pilot whale (*Globicephala macrorhynchus*) in Massachusetts, USA: 1980 and 2011. Aquat. Mamm. 42(3):357–362.
- Rone, B.K. and R.M. Pace, III. 2012. A simple photograph-based approach for discriminating between free-ranging long-finned (*Globicephala melas*) and short-finned (*G. macrorhynchus*) pilot whales off the east coast of the United States. Mar. Mamm. Sci. 28(2):254–275.
- Schwacke, L.H., E.O. Voit, L.J. Hansen, R.S. Wells, G.B. Mitchum, A.A. Hohn and P.A. Fair. 2002. Probabilistic risk assessment of reproductive effects of polychlorinated biphenyls on bottlenose dolphins (*Tursiops truncatus*) from the southeast United States coast. Env. Toxic. Chem. 21(12):2752–2764.
- Taruski, A.G., C.E. Olney and H.E. Winn. 1975. Chlorinated hydrocarbons in cetaceans. J. Fish. Res. Board Can 32(11):2205–2209.
- Thomas, L., J.L. Laake, E. Rexstad, S. Strindberg, F.F.C. Marques, S.T. Buckland, D.L. Borchers, D.R. Anderson, K.P. Burnham, M.L. Burt, S.L. Hedley, J.H. Pollard, J.R.B. Bishop and T.A. Marques. 2009. Distance 6.0. Release 2. [Internet]. University of St. Andrews (UK): Research Unit for Wildlife Population Assessment. Available from: <http://distancesampling.org/Distance/>.
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12, 93 pp. Available from: [http://www.nmfs.noaa.gov/pr/pdfs/sars/gamms\\_report.pdf](http://www.nmfs.noaa.gov/pr/pdfs/sars/gamms_report.pdf)
- Weisbrod, A.V., D. Shea, M.J. Moore and J.J. Stegeman. 2000. Bioaccumulation patterns of polychlorinated biphenyls and chlorinated pesticides in northwest Atlantic pilot whales. Environ. Toxicol. Chem. 19(3):667–677.
- Wells, R.S., J.B. Allen, G. Lovewell, J. Gorzelany, R.E. Delynn, D.A. Fauquier and N.B. Barros. 2015. Carcass-recovery rates for resident bottlenose dolphins in Sarasota Bay, Florida. Mar. Mamm. Sci. 31(1):355–368.
- Wells, R.S., E.M. Fougères, A.G. Cooper, R.O. Stevens, M. Brodsky, R. Lingenfelser, C. Dold, and D.C. Douglas. 2013. Movements and dive patterns of short-finned pilot whales (*Globicephala macrorhynchus*) released from a mass stranding in the Florida Keys. Aquat. Mamm. 39(1): 61–72.